# Valued Component – Other Mammals (Terrestrial) (Update November 2004)

# STATE OF KNOWLEDGE - WHAT IS HAPPENING?

A very brief overview of the state of knowledge with respect to terrestrial wildlife other than caribou and moose in the NWT is presented below. This overview is preliminary and not intended to be exhaustive.

- What are the baseline conditions with respect to terrestrial wildlife?
- There are 63 species of terrestrial mammals in the NWT. Many species are important to residents as a significant source of food, and as a source of income. Overall most species in the NWT are in good condition. Population numbers for individual species vary, and many factors influence the abundance and distribution of terrestrial wildlife. Major species are described below.

#### Wood Bison

Wood bison are found in four main areas south of the treeline in the NWT including the Liard and Nahanni River valleys, the Mackenzie bison range, Wood Buffalo National Park and the Slave River Lowlands. The Mackenzie and Nahanni bison herds are estimated at approximately 2000 and 350 animals respectively. Both herds are a result of conservation efforts to establish healthy bison populations that are not infected with bovine tuberculosis and brucellosis. Conversely, bison in Wood Buffalo National Park (ca. 4000 animals) and the adjacent Slave River Lowlands (ca. 500 animals) are infected with tuberculosis and brucellosis. The Bison Control Area (BCA) was created to prevent contact between the diseased and healthy bison populations; all bison in the BCA are presumed to be disease carriers and must be removed for testing. Ongoing surveillance of the Bison Control Area is important for minimizing the risk of disease transmission to healthy bison. Largely due to the disease issue, wood bison are listed as a threatened species in Canada, and considered at risk in the

#### KEY MONITORING INDICATORS

Population size and trendAge structureNumber harvestedDistributionHabitat qualityFood availability

Reproductive rates Presence of contaminants

Presence of diseases and parasites

Northwest Territories. The Mackenzie herd is an important conservation herd because it is the world's largest free ranging, wood bison herd. Another important conservation herd is the Hook Lake Wood Bison Recovery Project in Fort Resolution. This project was initiated in 1996, and has resulted in the establishment of a genetically important and disease-free captive breeding herd that will be used for national wood bison conservation efforts. Key indicators for monitoring wood bison herds are presence/absence of disease and population size.

#### Muskox

• Muskoxen are found throughout the tundra region and within the northern boreal habitat of the NWT. The muskox population on Banks Island (in the Inuvialuit Settlement Region) has greatly increased since the 1980s to a high of 69,000 in 2002 (after being nearly extirpated in the early 1900s). Substantial numbers of muskox are found in the area north of Great Bear Lake up to the arctic coast. Key indicators are population size, distribution and the presence of disease.

# Grizzly Bear

• Grizzly bear are found predominantly north of the treeline and in the Mackenzie Mountains. Population size in the Northwest Territories is estimated at 5100 bears. The grizzly is a species of special concern in Canada, and considered sensitive in the Northwest Territories. Monitoring by radio collar in the central barrens has shown that grizzlies can have large home ranges (averaging 7200 km² for males and 2100 km² for females).

Research is also taking place in the ISR. The population of grizzly bears is believed to be stable. Key indicators are population size, reproductive rates, habitat quality, and number harvested.

#### Black Bear

• Black bear are relatively abundant below the treeline in the NWT. The population is conservatively estimated at 10000, and believed to be healthy throughout its range. A key indicator is population size and trend, and number harvested.

#### Polar Bear

• Polar bear inhabit the coast lines of the Inuvialuit Settlement Region and are considered a vulnerable species in Canada. Populations are estimated at 3200 bears, divided into 3 subpopulations. Two sub-populations are considered stable while the other (shared with Nunavut) is recovering. Key monitoring indicators include presence of contaminants, reproductive rates, habitat quality, and number harvested.

# Lynx

• Lynx are found mainly below the treeline in the NWT. Their population mimics the 10-year population cycle of snowshoe hare (lagging behind 1 or 2 years). Peaks and lows can be predicted using snowshoe hare survey data. Populations in the western Northwest Territories appear to be increasing. Lynx populations are considered to be healthy. Key indicators are numbers harvested and food availability (snowshoe hare).

#### Fox

• Red fox and arctic fox are abundant in the NWT. Red fox are more abundant in the boreal forest while arctic fox thrive on the tundra. Key indicators are food availability, presence of disease such as rabies and number harvested.

## Wolf

• Wolves are found throughout the NWT. Tundra and timber (boreal) populations are considered to be healthy. Arctic wolves are found throughout the Arctic Archipelago and populations on Banks,

Melville and Victoria Islands are considered healthy. Key indicators are food availability, habitat quality, population size and pup survival, and number harvested.

#### Wolverine

 Wolverine population sizes are not known. They are a species of special concern in Canada, but considered secure in the Northwest Territories. Key indicators are habitat quality, reproductive rates and number harvested.

# Dall's Sheep

• Dall's sheep are found in the Mackenzie and Richardson Mountains along the Yukon border. Populations have been estimated at 14,000 to 26,000 based on a compilation of surveys of various mountain blocks. Dall's sheep populations are believed to be stable. Key indicators are population size, composition, reproduction and recruitment rates, harvest data, and the presence of parasites and disease.

#### Mountain Goat

• Mountain goats are a species that may be at risk in the Northwest Territories and are at the northern limit of their range. They are found only in the southern Mackenzie Mountains and have a limited distribution. Their population is estimated at 600 to 1000 and population trends are unknown. However, recent evidence in the Dehcho suggests that goat numbers and the extent of their range may be increasing. Key indicators are population size, distribution and habitat quality.

#### Beaver

Beaver are found primarily below the treeline in the NWT. They
have stable populations and are increasing in some regions. Beaver
are vulnerable to water level fluctuations and to localized over
harvest. Key indicators are numbers harvested and the density of
active lodges.

#### Marten

• Marten are found throughout forested areas of the NWT. Population cycles are largely influenced by prey availability, with numbers estimated at 40000 to 400000 depending on the stage in the cycle. Marten are harvested for their pelts. Populations are thought to be stable, but habitat loss and potential overharvesting are concerns. Key indicators are numbers harvested, age structure and ratio of females to males in the harvest, and food availability (index of small mammal abundance).

#### Mink

• Mink are found throughout the NWT, primarily in aquatic habitats. Their populations are believed to be stable, but are vulnerable to water pollution and localized overtrapping. They are considered to be an indicator of ecosystem health as they can bioaccumulate environmental pollutants. Key indicators are presence of contaminants, population size, and number harvested.

#### Small Mammals

- Small mammals are distributed widely throughout the NWT. They include snowshoe hare, arctic hare, porcupine and different species of weasels, squirrels, lemmings, mice, shrews and voles. Key indicators are indices of relative abundance (small mammal trapping, hare pellet counts and snow track counts).
- What are the levels and trends of contaminants in terrestrial wildlife?
- Contaminant levels in terrestrial wildlife are generally low in the NWT, and are not a concern from animal health or human food consumption perspectives.

# **CURRENT MONITORING**

Ongoing monitoring programs with respect to terrestrial wildlife in the NWT are found below.

# Overall monitoring

► NWT species status rank infobase (Government of the Northwest Territories since 2000).

This infobase monitors the general status of species in the Northwest Territories. It is a significant source of information for assessing future monitoring of terrestrial wildlife. The infobase is a searchable catalogue of information used to rank the status of species, thereby prioritizing them for further assessment and monitoring. The following biological indicators are used to rank species status: population size, number of occurrences, distribution, trend in population, trend in distribution, threats to population and threats to habitat. The general status ranks will be published every 5 years (see references for 2000 Status Ranks report).

 West Kitikmeot Slave Study 1996 –2001 (West Kitikmeot Slave Study Society)

WKSS was a trans-boundary program with GNWT, Federal, Nunavut and mining industry funding – focused on scientific and traditional knowledge studies in the Slave Geological Province area to gather baseline information for species such as caribou, grizzly bear, wolves and wolverine, as well as other ecological information. See Appendix A for more information on the WKSS.

► Traditional ecological knowledge research in the Kache Kue study region (Lutsel K'e Dene First Nation 1998-2001)

Traditional knowledge of Chipewyan elders will be recorded about the Kache Kue study region. The main focus is on the

habitat of key species and potential effects of resource development in the Slave Geological Province. Information was synthesized onto maps and indicators for ecosystem health were developed. Funding was provided through the WKSS.

- ► Northwest Territories wildlife disease and parasite monitoring (Government of the Northwest Territories since 1990)
- Northwest Territories commercial wildlife harvest monitoring (Government of the Northwest Territories since 1993) Monitors the commercial harvest of wildlife. Data collected includes the species and number harvested, date and location.
- Monitoring of non-resident and non-resident alien harvest from the Mackenzie Mountains (Government of the Northwest Territories since 1965)

Documents sport hunt harvest of big game species in the Mackenzie Mountains.

- ► Northwest Territories fur harvest monitoring (Government of the Northwest Territories since 1957)
  - Monitors the harvest of furbearers. Data collected includes the number of pelts of each species sold per trapper.
- Northwest Territories resident sport harvest monitoring (Government of the Northwest Territories since 1981)
   Monitors sport harvesting activities. Data collected includes species harvested, date and location by hunter.
- ► Gwich'in Settlement Area harvest study (Gwich'in Renewable Resource Board since 1995)
- ➤ Sahtu Settlement Area harvest study (Sahtu Renewable Resources Board 1998 2003)

- ▶ Inuvialuit Settlement Region harvest study (Inuvialuit Joint Secretariat 1986 2003)
- ► Harvest Data collection in Paulatuk, Holman, Sachs Harbour since 2001 (DFO, GNWT, and CWS)
- ► Harvest Data Collection in Aklavik (YTG)
- Wildlife Effects Monitoring Program (BHP Diamonds Inc. since 1994).

The program monitors caribou, grizzly bear, wolves, wolverines and foxes that pass through or live in the BHP claim block and may be affected by the EKATI diamond mine.

- ► Wildlife Monitoring Program (Diavik Diamond Mines Inc. since 1999)
- ▶ Wildlife/habitat monitoring at Snap Lake (DeBeers since 2002)
- ► Enbridge pipeline wildlife observation (Government of the Northwest Territories since 1999)
- ► Arctic Borderlands Ecological Knowledge (Cooperative)

#### Wood Bison

 Recovery of wood bison – ongoing population and disease monitoring (Government of the Northwest Territories)

#### Muskox

- Muskox mainland population surveys ISR / Sahtu (Government of the Northwest Territories)
- Muskox island population surveys ISR includes Banks and NW Victoria Islands. (Government of the Northwest Territories)

Muskox population surveys - Yukon (Parks Canada/YTG)

► Information on muskox health gathered during commercial harvest and urine/fecal collections (Government of the Northwest Territories)

# **Grizzly Bear**

- ► Continued monitoring of impacts from diamond mines in the Slave Geological Province (Government of the Northwest Territories since 2000)
- ► Grizzly bear harvest and occurrence databases, (Government of the NWT since 1977)

#### Black Bear

- ▶ Populations are monitored using harvest statistics as an indicator of population health.
- ► Monitoring of black bear diseases and parasites in the Dehcho. (Government of the Northwest Territories since 2002)

#### Polar Bear

- ► Polar bear harvest database (Government of the Northwest Territories since 1979)
- ► Monitoring and research studies on population boundaries and abundance through tagging programs (Canadian Wildlife Service and Government of the Northwest Territories)

#### Lynx

► Lynx population monitoring of harvest levels and pelt length measurements from pelts turned in for auction (Government of the Northwest Territories since 1985)

#### Fox

 Populations are monitored using harvest statistics and disease monitoring for rabies (Government of the Northwest Territories since 1957)

#### Wolf

- ► Wolf dens and pup counts on the central barrens (Government of the Northwest Territories since 1972)
- ► Wolf harvest monitoring the on Banks Island (Sachs Harbour) and NW Victoria Island (Holman) (Government of the Northwest Territories since 1990)
- ▶ Diet and morphometric measurements of Arctic wolves 1993-2002 (Government of the Northwest Territories)

#### Wolverine

- ► Populations are monitored using harvest statistics. (Government of the Northwest Territories since 1957)
- ► NWT wide wolverine carcass collection to improve monitoring of harvest patterns and health, age and sex composition (Government of the Northwest Territories starts in 2005)

# Dall's sheep

- Dall's sheep population size monitoring for Mackenzie Mountains, North Richardson Mountains, Canyon Creek subpopulation, Mount Cronin sub-population - Southern Richardson Mountains (Government of the Northwest Territories since 1970's)
- ▶ Parasite and disease studies of Dall's sheep in the Mackenzie Mountains (Government of the Northwest Territories since 1997)
- ► Population size monitoring in the Richardson Mountains (Government of the Northwest Territories)
- ► Monitoring growth rate pattern of harvested Dall's sheep rams in the Dehcho (Government of the Northwest Territories since 2002)

#### Mountain Goat

- ► Monitoring hunter reported observations of sex ratios in Mackenzie Mountains (Government of the Northwest Territories since 1995)
- ► Collecting DNA samples from goats harvested in the Dehcho, (Government of the Northwest Territories since 2002)

#### Beaver

► Beaver population monitoring - Sahtu (Government of the Northwest Territories since 1989)

#### Marten

 Marten carcass collection in select areas (for sex, age, population composition) (Government of the Northwest Territories)

#### Small Mammals

- Monitoring abundance of small mammals (Government of the Northwest Territories since 1990)
  - This program is designed to monitor the populations of small mammals across the Northwest Territories. Estimates of population size are based on the number of animals caught using snap and live traps.
- ➤ Studies of environmental effects of disturbances in the subarctic (SEEDS) Small mammals (University of Alberta since 1985)

  The program was established to investigate impacts of various disturbances associated with simulated transport corridors within upland subarctic ecosystems. The research site is 10 km north of Tulita, Northwest Territories. In 1995 a forest fire burned much of the site. Efforts to monitor post-fire ecosystem recovery continued until 2000. The microclimate installations have been removed but plans to continue small mammal and vegetation monitoring are being pursued.
- Snowshoe hare monitoring (Government of the Northwest Territories since 1989)

#### GAPS AND RECOMMENDATIONS FOR MONITORING

# Monitoring gaps and recommendations for future monitoring under the NWT Cumulative Impact Monitoring Program.

Given the range of species under consideration, decisions on monitoring gaps still requires further review amongst biologists, comanagement boards, and government agencies. Although few specific monitoring priorities have been established, several broad issues and recommendations are provided below:

- 1. Improve baseline information on wildlife distribution and abundance, and understanding of natural variability in populations.
- 2. Insufficient attention to cumulative effects relative to oil and gas exploration, pipeline construction, mining activity and anticipated increase in level of development activity in the north.
  - a) Require better habitat and land-use mapping and understanding of how wildlife utilize available habitat.
  - Requirement for standardized, broad multi-species survey techniques, collaborative monitoring efforts and data sharing
  - Absence of information to discriminate between effects of environmental variation and human activities.
- 3. There is uncertainty about the potential impact of global warming on northern wildlife species.

# REFERENCES

Relevant monitoring reports, past monitoring programs, research documents, and scientific publications are found below.

#### **OVERALL**

Carriere S. and J. Lange. 2002. **Detailed Proceedures Manual: General Status Ranks of Wild Species in the Northwesst Territories.** Manuscript Report 143. GNWT, 61 pp.

Dene Cultural Institute. Fort Good Hope - Colville Lake traditional environmental knowledge pilot project (1989-1993).

Elkin, B.T. (1999). **Community-based monitoring of abnormalities** *in wildlife*. In: J. Jensen (ed.), Synopsis of Research Conducted Under the 1997/98 Northern Contaminants Program. pp. 123-126. Environmental Studies No. 75, Indian and Northern Affairs Canada. 434 pp.

This program was funded through the Northern Contaminants Program from 1997 to 2000. It was a pilot study for a community-based monitoring system, to provide "early warning" of abnormalities/changes in wildlife health. The project focused on the communities of Fort Good Hope, Fort Resolution and Kugluktuk (Nunavut), and sought to integrate traditional knowledge with western science.

Government of the Northwest Territories (2000). **NWT Species 2000: General status ranks of wild species in the Northwest Territories**. (see Current Monitoring for description).

Government of the Northwest Territories. **Norman Wells wildlife monitoring program** (1983-1986).

Nagy, J. A., T. Creighton, W. Wright, T. Slack, M. Branigan, and D. Auriat. 2003. **Wildlife research and information gaps workshop**,

draft workshop summary for review by communities. Western NWT Biophysical Study, Department of Resources, Wildlife, and Economic Development, Inuvik and Yellowknife. 72 pp.

Penner and Associates Ltd. (1997). Wildlife baseline studies progress report, 1995 and 1996 investigations, Diavik Diamond Mines Inc. Project, Lac de Gras, NWT: Draft report. In: Diavik Diamond Mines Inc. 1998. Project description submission. Penner and Associates Ltd.

Kutz, S.J., Hoberg, E.P., Nagy, J.A., Polley, L. and Elkin, B. 2004. "Emerging" parasitic infections in arctic ungulates. Integrated Comp. Biol. 44: 109-118.

## **WOOD BISON**

Bidwell W.A., Nishi J.S., Ellsworth T.R. 2004. **Bison Control Area** program Annual Report of Survey Activities Dec 2003 - April 2004 Government of the Northwest Territories Manuscript Report No. 156. 50pp

Gates, C.C. and N.C. Larter. 1990. **Growth and dispersal of an erupting large herbivore population in Northern Canada: the Mackenzie wood bison (Bison bison athabascae).** Arctic. 43: 231-238.

Gates, C.C., N.C. Larter and P.E. Komers. 1989. **Size and composition of the Mackenzie wood bison population in 1989**. G.N.W.T Renewable Resources File Rep. No. 93. 24pp.

Gates, C.C., R.O. Stephenson, H.W. Reynolds, C.G. van Zyll de Jong, H. Schwantje, M. Hoefs, J. Nishi, N. Cool, J. Chisholm, A. James, and B. Koonz. 2001. **National Recovery Plan for the Wood Bison (Bison bison athabascae).** National Recovery Plan No. 21. Recovery of Nationally Endangered Wildlife (RENEW). Ottawa, Ontario. 50 pp.

Government of the Northwest Territories (2000). **Species at risk** fact sheet - NWT wood bison (Bison bison athabascae)

Canadian Status: Threatened. From the Wildlife & Fisheries website, <a href="http://www.nwtwildlife.rwed.gov.nt.ca">http://www.nwtwildlife.rwed.gov.nt.ca</a>.

Komers, P. F. Messier, and C.C. Gates. 1993. **Group structure in wood bison: nutritional and reproductive determinants.**Canadian Journal of Zoology. 71: 1367-1371.

Larter, N.C. and C.C. Gates. 1990. Home ranges of wood bison in an expanding population. Journal of Mammology. 71: 604-607

Larter, N.C. and C.C. Gates. 1991. **Diet and habitat selection of wood bison in relation to seasonal changes in forage quantity and quality.** Canadian Journal of Zoology. 69: 2677-2685.

Larter, N.C. and C.C. Gates. 1994. **Home ranges of wood bison:** the effects of age, sex, and forage availability. J. Mamm. 75: 142-149.

Larter, N.C., A.R.E. Sinclair and C.C. Gates. 1993. **Dynamics of the Mackenzie wood bison population**. Proceedings of the 2<sup>nd</sup> North American Public Bison Herds Symposium, pp. 260-270.

Larter, N.C., A.R.E. Sinclair and C.C. Gates. 1994. **The response** of predators to an erupting Bison (Bison bison athabascae) population. Can. Field-Nat. 108: 318-327.

Larter N.C., Sinclair A.R.E., Ellsworth T., Nishi J. and Gates C.C. 2000. **Dynamics of reintroduction in an indigenous large ungulate: the wood bison of northern Canada.** Animal Conservation 4: 299–309

Larter, N.C., J.S. Nishi, T. Ellsworth, D. Johnson, G. More and D.G. Allaire. 2003. **Observations of wood bison swimming across the Liard River, Northwest Territories, Canada**. Arctic 56: 408-412.

Melton, D.A., N.C. Larter, C.C. Gates and J.A. Virgl. 1989. The influence of rut and environmental factors on the behaviour of wood bison. Acta Theriol. 34: 179-193.

Melton, D.A., C.C. Gates, N.C. Larter and J.A. Virgl. 1990. Foraging behaviour of wood bison in an expanding population. <u>In:</u>
Transactions of the XIX IUGB Congress Volume 1: Population Dynamics (S. Myrberget ed.). Norwegian Institute for Nature Research, Trondheim pp. 44-47.

Nishi, J.S., 2002. Surveillance Activities Under the Northwest Territories Bison Control Area Program (1987-2000). Department of Resources, Wildlife and Economic Development, Government of the Northwest Territories, Manuscript Report No. 145. 32p.

Nishi, J.S. B.T. Elkin, and T.R. Ellsworth. 2002. The Hook Lake Wood Bison Recovery Project: Can a disease-free captive wood bison herd be recovered from a wild population infected with bovine tuberculosis and brucellosis? Annals of the New York Academy of Sciences. 969: 229-235.

Wilson, G.A. and C. Strobeck. 1999. **Genetic variation within and relatedness among wood and plains bison populations.** Genome 42:483-496.

#### MUSKOX

Case, R.L. and K.G. Poole (1985). **Distribution, abundance and composition of muskoxen north of Great Bear Lake, March 1983**. Government of the Northwest Territories. 48pp.

Fournier, B. and A. Gunn (1997). **Muskox numbers and** distribution in the Northwest Territories, 1997. Government of the Northwest Territories. 55pp.

Graf, R. and C. Shank (1989). **Abundance and distribution of muskoxen near Artillery Lake, NWT, March 1989.** Government of the Northwest Territories. 19pp.

Hoberg, E.P., Kutz S.J., Nagy J., Jenkins E., Elkin B., Branigan, M., and Cooley D. 2002. **Protostrongylus stilesi (Nematoda: Protostrongylidae): Ecological Isolation and Putative Host-**

**Switching Between Dall's Sheep and Muskoxen in a Contact Zone** Comparative Parasitology 69(1):1-9.

McLean, B.D. (1992). An aerial survey of muskoxen north of Great Bear lake, August 1987. Government of the Northwest Territories. 19pp.

McLean, B.D. and P. Fraser (1992). Abundance and distribution of Peary caribou and muskoxen on Banks Island, NWT, June 1989. Government of the Northwest Territories. 28pp.

Larter, N.C. 1999. An aerial survey of muskoxen in the Paulatuk area of the Inuvialuit Settlement Region, March 1997. G.N.W.T. Resources, Wildlife & Economic Development Manuscript Rep. No. 116. 27pp.

Larter, N.C. and J.A. Nagy. 1997. **Peary caribou, muskoxen, and Banks Island forage: assessing seasonal diet similarities**. Rangifer 17: 9-16.

Larter, N.C. and J.A. Nagy. 1999a. **Muskox mortality survey, Banks Island August, 1996.** G.N.W.T. Resources, Wildlife & Economic Development Manuscript Rep. No. 117. 13pp.

Larter, N.C. and J.A. Nagy. 1999b. **Sex and age classification surveys of muskoxen on Banks Island, 1985-1998: a review**. G.N.W.T. Resources, Wildlife & Economic Development Manuscript Rep. No. 113. 32pp.

Larter, N.C. and J.A. Nagy. 2001a. **Overwinter changes in the urine chemistry of muskoxen from Banks Island.** J. Wildl. Manage. 65: 226-234.

Larter, N.C. and J.A. Nagy. 2001b. Variation between snow conditions at Peary caribou and muskox feeding sites and elsewhere in foraging habitats on Banks Island in the Canadian High Arctic. Arct. Antarct. Alp. Res. 33: 123-130.

Larter, N.C. and J.A. Nagy. 200lc. Calf production, calf survival, and recruitment of muskoxen on Banks Island during a period of changing population density from 1986-1999. Arctic 54: 394-406.

Larter, N.C. and J.A. Nagy. 2003 Wet sedge meadow habitat of southern Banks Island excluded from grazing by large herbivores for five years: effects on aboveground standing crop. G.N.W.T. Resources, Wildlife & Economic Development Manuscript Rep. No. 150. 32pp.

Larter, N.C. and J.A. Nagy. 2004 **Seasonal changes in the** composition of the diets of Peary caribou and muskoxen on **Banks Island.** Polar Research in press

Larter, N.C., J.A. Nagy and D.S. Hik. 2002. **Does seasonal** variation in forage quality influence the potential for resource competition between muskoxen and Peary caribou on Banks Island? Rangifer 22: 143-153.

Nagy, J.A. N.C. Larter and V.P. Fraser. 1996. **Population** demography of Peary caribou and muskox on Banks Island, N.W.T., 1982-1992. Rangifer Special Issue 9: 213-222.

Kutz, S., Garde, E., Veitch, A., Nagy, J., Ghandi, F. and Polley, L. 2004. Muskox lungworm (Umingmakstrongylus pallikuukensis) does not establish in experimentally exposed thinhorn sheep (Ovis dalli). J. Wildl. Dis. 40: 197-204.

Barboza, P.S., Rombacj, E.P., Blake, J.E. and Nagy, J.A. 2003. **Copper status of muskoxen: a comparison of wild and captve populations.** J. Wildl. Dis. 39: 610-619.

#### **GRIZZLY BEAR**

Government of the Northwest Territories (2000). **Species at risk** fact sheet - NWT grizzly bear (Ursus arctos horribilis) Canadian Status: Special concern. From the Wildlife & Fisheries website, http://www.nwtwildlife.rwed.gov.nt.ca.

Government of the Northwest Territories. **Bear tracks: A newsletter on grizzly bear studies in the Central Arctic, NWT, Canada**. (1996-2000).

Eight issues have been published to date. Issues are found at <a href="http://www.nwtwildlife.rwed.gov.nt.ca/Publications/beartracks/">http://www.nwtwildlife.rwed.gov.nt.ca/Publications/beartracks/</a>.

Government of the Northwest Territories. Grizzly bear (Ursus Actos) studies in the Northwest Territories: Nutritional component. Final Report to the West Kitikmeot/Slave Study (released June 14, 1999)

Government of the Northwest Territories. **Grizzly bear productivity** in the Richardson Mountains (1993-2000).

McLoughlin, P.D., R.L. Case, R.J. Gau, S.F. Ferguson and F. Messier (1999). Annual and seasonal movement patterns of barren-ground grizzly bears in the central Northwest Territories. Ursus Vol. 11.

McLoughlin, P.D., F. Messier, R.L. Case, R.J. Gau, R. Mulders and H.D. Cluff (1999). The spatial organization and habitat selection patterns of barren-ground grizzly bears (Ursus arctos) in the Northwest Territories and Nunavut. (1995-1999) Final report to the West Kitikmeot Slave Study Society. Yellowknife, NT. 54pp.

McLoughlin, P.D., R.L. Case, R.J. Gau, H.D. Cluff, R. Mulders and F. Messier (in review). Effect of temporal and spatial differences in habitat on home ranges of barren-ground grizzly bears in the central Arctic. Canadian Journal of Zoology.

McLoughlin, P.D., H.D. Cluff, R.J. Gau, R. Mulders, R.L. Case and F. Messier (2002). Hierarchical habitat selection patterns of barrenground grizzly bears in Nunavut and the Northwest Territories, Canada. Oecologica 132:102-108.

Nagy, J.A. and M. Branigan. 1998. Co-management plan for grizzly bears in the Inuvialuit Settlement Region, Yukon

Territory and Northwest Territories, with work plans for 1997/1998 to 2001/2002. 63 pp.

Strom, K.B. and P.D. McLoughlin (1998). Habitat effectiveness model for grizzly bears in the Lac de Gras region, Northwest Territories, Canada. Technical report for the Diavik Diamonds Project Environmental Effects Report for Wildlife. Calgary, AB.

#### WOLF

Carmichael, L., J.A. Nagy, N.C. Larter and C. Strobeck. 2001. **Prey specialization may influence patterns of gene flow in wolves of the Canadian Northwest.** Molecular Ecol. 10: 2787-2798.

Cluff, H.D., L.R. Walton and P.C. Paquet (2001). Movements and habitat use of wolves denning in the central Arctic, Northwest Territories and Nunavut. (1996-1999) Final Report to the West Kitikmeot Slave Study Society, Yellowknife, NT.

Frame, P.F. 2004. Effects of caribou movements and anthropogenic development on tundra denning wolves in the central Canadian arctic. M.Sc. thesis, University of Alberta.

Frame, P.F., H.D. Cluff, and D.S. Hik. (in review). Effects of seasonal caribou movements and human development on reproductive success of tundra denning wolves in the central Canadian arctic. Submitted to Canadian Journal of Zoology.

Frame, P.F., H.D. Cluff, and D.S. Hik. (in review). Response of wolves to experimental disturbance at home sites in the central Canadian arctic. Submitted to Conservation Biology.

Frame P.F, D.S. Hik, H.D.Cluff and P.C.Paquet. 2004. Long Foraging Movement of a Denning Tundra Wolf. Arctic 57:196-203

Cluff D., and P. Paquet. 2003. Large Carnivores as Umbrellas for Reserve Design and Selection in the North. 2003. Canadian Council on Ecological Areas

*Musiani, M. 2003.* Conservation biology and management of wolves and wolf-human conflicts in western North America. Ph.D. dissertation, University of Calgary.

Nagy, J.A. and N.C. Larter. 2000. Status and diet of arctic wolves (Canis lupus arctos) in the Inuvialuit Settlement Region, Arctic Canada. Abstract only. Beyond 2000 Realities of Global Wolf Restoration. Univ. Michigan, Duluth, MN pg 91.

Government of the Northwest Territories. Wolf notes: A newsletter on wolf studies in the Central Arctic, NWT, Canada. (1997-2000). Five issues have been published to date. Issues are found at <a href="http://www.nwtwildlife.rwed.gov.nt.ca/Publications/wolfnotes/">http://www.nwtwildlife.rwed.gov.nt.ca/Publications/wolfnotes/</a>

Government of the Northwest Territories. Organochloride, heavy metal and radionuclide contaminant transfer through the lichencaribou-wolf food chain (1992-1996).

Walton, L.R. (2000). **Investigation into the movements of migratory wolves in the central Canadian Arctic.** MSc Thesis, University of Saskatchewan, Saskatoon, Canada. 59pp.

Walton, L.R., H.D. Cluff, P.C. Paquet and M.A. Ramsay (2001). **Movement patterns of barren-ground wolves in the central Canadian arctic**. Journal of Mammalogy 82(3): in press.

#### WOLVERINE

Mulders, R. (2000) Wolverine ecology, distribution and productivity in the Slave Geological Province (1995-1999). West Kitikmeot Slave Study, 90 pgs.

Government of the Northwest Territories (2000). **Species at risk** fact sheet - NWT wolverine (Gulo gulo) Canadian Status: Special concern. From the Wildlife & Fisheries website, http://www.nwtwildlife.rwed.gov.nt.ca.

#### DALL'S SHEEP

Case, R. (1989). **Distribution and abundance of Dall's sheep in the southern Mackenzie Mountains, NWT.** Government of the Northwest Territories. 39pp.

Ferguson, R.S., K.D. Davidge and R.S. Douglas (1985). **Distribution and abundance of Dall's sheep near Jackfish River, Northwest Territories in June 1984**. Government of the Northwest Territories. 20pp.

Kutz, S.J., A.M. Veitch, E.P. Hoberg, B.T. Elkin, E.J. Jenkins and L.. Polley (2000). New host and geographic records for Protostrongylus stilesi and Parelaphostrongylus odocolei in Dall's sheep (Ovis dalli dalli) from the Mackenzie Mountains, Northwest Territories, Canada. Journal of Wildlife Diseases. 37: 761-744.

Veitch, A.M., E. Simmons, J. Adamczewski and R. Popko (2000). Status, harvest and co-management of Dall's sheep in the Mackenzie Mountains, NWT 1997. Northern Wild Sheep and Goat Council 11 (1998). 11: 134-153.

Larter, N.C. and D.G. Allaire. 2004. **Mackenzie mountain non-resident and non-resident alien hunter harvest summary 2003.** G.N.W.T. Resources, Wildlife & Economic Development Manuscript Rep. No. 154. 46pp.

#### **MOUNTAIN GOAT**

Veitch, A., E. Simmons, M. Promislow, D. Tate, M. Swallow, and R. Popko 2002. **The status of mountain goats in Canada's Northwest Territories.** Proceedings of the 13<sup>th</sup> Northern Wild Sheep and Goat Council: 49-61.

Simmons, N.M. 1982a. Seasonal range of Dall's sheep, Mackenzie Mountains, Northwest Territories. Arctic 35: 512-518. Simmons, N.M. 1982b. **Seasonal distribution of Dall's sheep in the Mackenzie Mountains, Northwest Territories**. NWT Wildl. Serv. Rep. No. 21 47pp.

Larter, N.C. and Allaire, D.G. in press. **Sheep surveys of the Liard Range, Nahanni Range and Ram Plateau in the Mackenzie Mountains, August 2003.** GNWT Ms. Rep.

Worley, K., Strobeck, C., Arthur, S., Carey, J., Schwantje, H. and Veitch, A. 2004. **Population genetic structure of North American thinhorn sheep (Ovis dalli)**. Molecular Ecol. 13: 2545-2556.

#### **MOUNTAIN GOAT**

Veitch, A., E. Simmons, M. Promislow, D. Tate, M. Swallow, and R. Popko (2002). **The status of mountain goats in Canada's Northwest Territories.** Proceedings of the 13<sup>th</sup> Northern Wild Sheep and Goat Council: 49-61.

Larter, N.C. 2004. Mountain goat survey, Flat River area, western Mackenzie Mountains, September 2004 G.N.W.T. Resources, Wildlife & Economic Development Manuscript Rep. No. 157, 16pp in press.

#### MINK

Poole, K.G., B.T. Elkin and R.W. Bethke (1995). **Environmental contaminants in wild mink in the Northwest Territories, Canada**. Science of the Total Environment, 160/161: 473-486.

This Government of the Northwest Territories program examined organochlorine and heavy metal residues in wild mink from the Northwest Territories from 1991 to 1995.

Poole, K.G., B.T. Elkin and R.W. Bethke (1998). **Organochlorine** and heavy metal contaminants in wild mink in western **Northwest Territories, Canada**. Archives of Environmental Contamination and Toxicology, 34:406-413.

#### BEAVER/MUSKRAT

Fort Resolution Environmental Working Committee and Indian and Northern Affairs Canada. **Metals and organic contaminants in beaver and muskrat in the Slave River Delta area, NWT**. Northern Contaminants Program (1998-1999).

Popko, R. A. and A. M. Veitch (1998) **An aerial survey for beavers** in the Sahtu Settlement Area, October 1997. Government of the Northwest Territories. 21 pp.

Popko, R. A., A. M. Veitch, and M. C. Promislow (2002). An aerial survey for beaver lodges in the Sahtu Settlement Area, September 2001. Government of the Northwest Territories. 25 pp.

#### SMALL MAMMALS

Krebs, J. K., A.J. Kenney, S. Gilbert, K Danell, A. Angerbjorn, S. Erlinge, R.G. Bromley, C. Shank and S. Carriere. 2002. **Syncrony in lemming and vole populations in the Canadian Arctic**. Can. J. Zool. 80: 1323-1333.

Shank, C. (1997). **The Northwest Territories small mammal survey: 1990-1996**. Government of the Northwest Territories. Manuscript #96.

Larter, N.C. 1998. Collared lemming abundance, diet, and morphometrics on Banks Island, 1993-1996. G.N.W.T. Resources, Wildlife & Economic Development Manuscript Rep. No. 107. 21pp.

Larter, N.C. 1999b. **Seasonal changes in arctic hare, Lepus arcticus, diet composition and differential digestibility**. Can. Field-Nat. 113: 481-486.